arXiv:1512.08686v1 [math.GN] 29 Dec 2015

SOME NEW QUESTIONS ON POINT-COUNTABLE COVERS AND SEQUENCE-COVERING MAPPINGS

SHOU LIN AND JING ZHANG

Abstract. In this survey, 37 questions on point-countable covers and sequence-covering mappings are listed. These questions are mainly related to the theory of generalized metric spaces, involving point-countable covers, sequence-covering mappings, images of metric spaces and hereditarily closure-preserving families.

1. Introduction

All the time, the problems in mathematics are a powerful driving force to lead the development of mathematics [26]. Since *Open Problems in Topology* [69] and *Open Problems in Topology II* [71] were published, general topology and related fields have obtained huge development [25, 27, 28]. The new version of the book *Point-countable Covers and Sequence-covering Mappings* [43] written by the listed author will be published soon. This book is devoted to the area of some classes of generalized metric spaces, which are defined by certain covers or networks. It systemically investigates the theory of spaces with point-countable covers and sequence-covering mappings on generalized metric spaces, including the celebrated works obtained by Chinese scholars in this field, such as point-countable covers, sequences of point-finite covers, hereditarily closure-preserving covers and star-countable covers. It is worth mentioning that there are 91 open problems were listed in this book.

In the preface to the first edition of this book [43], Arhangel'skiĭ stated: "Though reading the monograph does not require much special background, the exposition in it goes far beyond the elementary level. It contains a rich collection of deep and beautiful results of highest professional level. The book not only brings the readers to the very first line of investigations in the theory of generalized metric spaces, but also contains many intriguing and important unsolved problems, some of them old and some new. $\cdots$ I would like to mention another joyful aspect of this monograph. Its appearance

2010 *Mathematics Subject Classification.* Primary 54E40; Secondary 54C10; 54E35; 54D70; 54-02.

Key words and phrases. Metric spaces; Point-countable families; Sequence-covering mappings; Hereditarily closure-preserving families; k-networks; cs-networks.

Supported in part by the NSFC (Nos. 11201414, 11471153) and the Training Programme Foundation for Excellent Youth Researching Talents of Minnan Normal University (No. MJ14001).

marks the success of a long period of development of general topology in China, it brings to the light important contributions to the mainstream of general topology made by a very creative group of Chinese mathematicians."

Because this book is published in Chinese, it is difficult to understand Chinese topologist' contributions and related problems in this field for the readers who are not proficient in Chinese. To provide convenience for the readers who are interested to continue in-depth study, we selects 37 questions from this book.

All spaces in this paper are assumed to be T_2, and all mappings are continuous and onto. First, recall some basic definitions.

Definition 1.1. Let X be a topological space, and $\mathscr{P}$ a family of subsets of X.

(1) $\mathscr{P}$ is called a *k-network* for X [70] if whenever $K \subset V$ with K compact and V open in X, there exists a finite subfamily $\mathscr{P}' \subset \mathscr{P}$ such that $K \subset \cup\mathscr{P}' \subset V$.
(2) $\mathscr{P}$ is called a *cs-network* for X [24] if, given a sequence $\{x_n\}$ converging to a point $x \in X$ and a neighborhood V of x in X, then $\{x\} \cup \{x_n : n \geq n_0\} \subset P \subset V$ for some $n_0 \in \mathbb{N}$ and some $P \in \mathscr{P}$.
(3) $\mathscr{P}$ is called a *cs*-network* for X [19] if whenever $\{x_n\}$ is a sequence converging to a point $x \in V$ with V open in X, then $\{x\} \cup \{x_{n_i} : i \in \mathbb{N}\} \subset P \subset V$ for some subsequence $\{x_{n_i}\}$ of $\{x_n\}$ and some $P \in \mathscr{P}$.
(4) $\mathscr{P}$ is called a *wcs*-network* for X [49] if whenever $\{x_n\}$ is a sequence converging to a point $x \in V$ with V open in X, then $\{x_{n_i} : i \in \mathbb{N}\} \subset P \subset V$ for some subsequence $\{x_{n_i}\}$ of $\{x_n\}$ and some $P \in \mathscr{P}$.

Definition 1.2. Let $\mathscr{P} = \bigcup_{x \in X} \mathscr{P}_x$ be a family of subsets of a topological space X satisfying as follows: (a) for each $x \in X$, $\mathscr{P}_x$ is a *network* of x in X, i.e., $\mathscr{P}_x \subset (\mathscr{P})_x$ and if $x \in G$ and G is open in X, then $P \subset G$ for some $P \in \mathscr{P}_x$; (b) if $U, V \in \mathscr{P}_x$, then $W \subset U \cap V$ for some $W \in \mathscr{P}_x$.

(1) $\mathscr{P}$ is an *sn-network* for X [39] if each element P of $\mathscr{P}_x$ is a sequential neighborhood of x for each $x \in X$, i.e., every sequence converging to the point x is eventually in P. The $\mathscr{P}_x$ is called an *sn*-network at x in X.
(2) $\mathscr{P}$ is called a *weak-base* for X [8] if for every $G \subset X$, the set G is open in X whenever for each $x \in G$ there exists a $P \in \mathscr{P}_x$ such that $P \subset G$. The $\mathscr{P}_x$ is called a *weak-base* at x in X.
(3) A space X is *snf-*(resp. *gf-*)*countable* [8, 40] if X has an *sn*-network (resp. a *weak-base*) $\mathscr{P}$ such that each $\mathscr{P}_x$ is countable.

The following relations hold for a topological space [43]:

(1) bases $\Rightarrow$ weak-bases $\Rightarrow$ *sn*-networks $\Rightarrow$ *cs*-networks $\Rightarrow$ *cs**-networks $\Rightarrow$ *wcs**-networks.
(2) bases $\Rightarrow$ *k*-networks $\Rightarrow$ *wcs**-networks.

Definition 1.3. Let X, Y be topological spaces, and $f : X \to Y$ a mapping.

(1) f is a *compact mapping* (resp. *s-mapping*) if each $f^{-1}(y)$ is compact (resp. separable) in X.
(2) f is a *boundary-compact mapping* (resp. *at most boundary-one mapping*) if each $\partial f^{-1}(y)$ is compact (resp. at most one point) in X.
(3) f is a *compact-covering mapping* [66] if each compact subset of Y is the image of some compact subset of X.
(4) f is a *sequence-covering mapping* [80] if whenever $\{y_n\}$ is a convergent sequence in Y, there exists a convergent sequence $\{x_n\}$ in X with each $x_n \in f^{-1}(y_n)$.
(5) f is a *pseudo-sequence-covering mapping* [23, 29] if whenever S is a convergent sequence containing its limit point in Y, there is a compact subset K in X such that $f(K) = S$.
(6) f is a *sequentially quotient mapping* [13] if whenever $\{y_n\}$ is a convergent sequence in Y, there exist a subsequence $\{y_{n_i}\}$ of $\{y_n\}$ and a convergent sequence $\{x_i\}$ in X such that each $x_i \in f^{-1}(y_{n_i})$.
(7) f is a 1-*sequence-covering mapping* [39] if for each $y \in Y$, there is an $x \in f^{-1}(y)$ such that whenever $\{y_n\}$ is a sequence converging to y in Y, there is a sequence $\{x_n\}$ converging to x in X with each $x_n \in f^{-1}(y_n)$.

The following is obvious:

(1) 1-sequence-covering mappings $\Rightarrow$ sequence-covering mappings $\Rightarrow$ pseudo-sequence-covering mappings, and sequentially quotinet mappings.
(2) boundary-compact closed mappings $\Rightarrow$ compact-covering mappings $\Rightarrow$ pseudo-sequence-covering mappings.

Readers may refer to R. Engelking's standard book *General Topology* [17] and G. Gruenhage's fine paper *Generalized Metric Spaces* [22] for unstated definitions and terminology.

2. Point-countable Covers

Spaces with a point-countable base can be considered as the most beautiful ones with a point-countable cover. Many results on spaces with a point-countable cover can be traced back to certain properties of metric spaces, in particular, spaces having a point-countable base. Every space with a point-countable base is meta-Lindelöf. Further,

(1) Every Fréchet space with a point-countable cs^*-network is meta-Lindelöf [23, 43].
(2) Every regular Fréchet space with a point-countable k-network is meta-Lindelöf [23].
(3) There exists a regular space with a point-countable weak-base which is not meta-Lindelöf [23].
(4) There exists a first-countable space having a σ-discrete k-network which is not meta-Lindelöf [43, Example 1.4.8].

Now, we introduce a concept as follows [43, Definition 2.1.15]. Let X be a topological space and $\mathscr{P}$ a family of subsets of X. $\mathscr{P}$ is called a *cs'-network* for X if whenever $\{x_n\}$ is a sequence in X converging to a point $x \in U$ with U open in X, there exist an $n \in \mathbb{N}$ and a $P \in \mathscr{P}$ such that $\{x, x_n\} \subset P \subset U$. Obviously, each cs^*-network is a cs'-network in a space.

Question 2.1. ([43, Question 2.1.24]) Is every Fréchet space with a point-countable cs'-network meta-Lindelöf?

Every k-space with a point-countable k-network is preserved by closed mappings [49, 79]. However, spaces with a point-countable k-network are not necessarily preserved by closed mappings [74]. As each compact subset of a space having a point-countable k-network is metrizable [23], we have the following question.

Question 2.2. ([43, Question 2.3.7]) Let X and Y be topological spaces, and $f : X \to Y$ a compact-covering and closed mapping. Does the space Y have a point-countable k-network if X has a point-countable k-network and each compact subset of Y is metrizable?

Let $\mathscr{P}$ be a family of subsets of a topological space X. $\mathscr{P}$ is called *compact-countable* if every compact subset of X meets at most countably many elements of $\mathscr{P}$.

Question 2.3. ([43, Question 2.3.16], [45]) Is every k-space with a compact-countable k-network preserved by closed mappings?

Michael, Nagami [67] posed the following famous problem: Is every quotient s-image of a metric space also a compact-covering quotient and s-image of a metric space? However, Chen gave a negative answer to this problem as the following two examples shows:

(1) there is a space which is a quotient compact image of a metric space, but not any compact-covering quotient and s-image of a metric space [15];
(2) there is a regular space which is a quotient s-image of a metric space, but not any compact-covering quotient and s-image of a metric space under the condition with a σ'-set [16].

In order to characterize compact-covering s-images of metric spaces, we recall the concept of cfp-networks.

A family $\mathscr{P}$ of subsets of a topological space X is called a *cfp-network* for X [87] if for every compact subset K of X and a neighborhood V of K in X there exists a finite subfamily $\mathscr{P}' \subset \mathscr{P}$ such that $\mathscr{P}'$ can be precisely refined by a finite cover of K consisting of closed subsets of K and $\cup\mathscr{P}' \subset V$. Obviously, every closed k-network is a cfp-network, and every cfp-network is a cs^*-network in a topological space.

Sequential spaces with a point-countable cs^*-network can be characterized as quotient s-images of metric spaces [83], and k-spaces with a point-countable cfp-network can be characterized as compact-covering quotient

and s-images of metric spaces [11, 87]. Chen's example shows that a sequential space with a point-countable cs^*-network may not have any point-countable cfp-network.

Question 2.4. ([43, Question 2.5.21])

(1) Does every sequential space with a compact-countable cs^*-network have a point-countable cfp-network?
(2) Is there a regular space X which has a point-countable cs^*-network but no any point-countable cfp-network under (ZFC)?

A regular space is metrizable if and only if it has a σ-compact-finite base [12]. Closed images of metric spaces can be characterized by regular Fréchet spaces with a σ-compact-finite k-network [56]. It remains an open problem [63] whether a regular space with a σ-compact-finite weak-base has a σ-locally finite weak-base. Banakh, Bogachev and Kolesnikov [10] proved that a separable regular space with a σ-compact-finite k-network has a countable k-network under (CH).

Question 2.5. ([43, Question 4.1.24]) Does every separable regular space with a σ-compact-finite weak-base have a countable weak-base under (ZFC)?

A compact closed mapping is called a *perfect mapping* [17]. A countable-to-one perfect image of a space with a countable weak-base is not necessarily gf-countable.

Question 2.6. ([43, Question 2.6.24], [88]) Is every space with a point-countable weak-base preserved by finite-to-one closed mappings?

Michael, Nagami [67] showed that each compact subset of a space with a point-countable base has a countable outer base. Let $\mathscr{P} = \bigcup_{x\in X} \mathscr{P}_x$ be a weak-base for a space X in Definition 1.2(2) and $A \subset X$. The family $\bigcup_{x\in A} \mathscr{P}_x$ is called an *outer weak-base* [54] of the set A in X.

Question 2.7. ([43, Question 2.7.20], [54]) Does every compact subset of a space with a point-countable weak-base have a countable outer weak-base?

Let $\mathscr{P}$ be a cover of a topological space X. $\mathscr{P}$ is called *point-regular* [1] if $x \in U$ and U is open in X, then $\{P \in (\mathscr{P})_x : P \not\subset U\}$ is finite.

(1) Spaces with a point-regular base can be characterized as (compact-covering) open and compact images of metric spaces [5].
(2) Spaces with a point-regular weak-base can be characterized as sequence-covering quotient and compact images of metric spaces [29].
(3) Spaces with a point-regular cs-network (resp. sn-network) can be characterized as sequence-covering compact images of metric spaces [52].
(4) Spaces with a point-regular cs^*-network can be characterized as pseudo-sequence-covering (and sequentially quotient) compact images of metric spaces [3].

Question 2.8. ([43, Question 3.4.20], [55]) How to characterize spaces with a point-regular k-network by certain images of metric spaces?

Shakhmatov [76] and Watson [86] constructed independently the following famous example. There exists a completely regular pseudo-compact non-compact space X having a point-countable base. Uspenskii [85] proved that pseudo-compact spaces with a σ-point-finite base is metrizable.

Question 2.9. ([43, Question 3.4.21]) Is every pseudo-compact space with a point-regular weak-base metrizable?

3. Sequence-covering mappings

A finite-to-one closed mapping defined on a metric space is not necessarily a sequence-covering mapping [80]. Closed mappings on regular spaces in which each point is a G_δ-set are sequentially quotient mappings [43, Lemma 2.3.3].

Question 3.1. ([43, Question 2.3.15], [41]) Is every closed mapping on spaces in which each point is a G_δ-set sequentially quotient?

Siwice [80] showed that every open mapping on a first-countable space is a sequence-covering mapping. In fact, every almost-open mapping on a first-countable space is a 1-sequence-covering mapping [42]. Let X and Y be topological spaces, and $f : X \to Y$ a mapping. f is called an *almost-open mapping* [6] if for each $y \in Y$, there is an $x \in f^{-1}(y)$ such that whenever U is a neighborhood of x in X, then $f(U)$ is a neighborhood of y in Y. There exists an open mapping on a Fréchet space such that it is not pseudo-sequence-covering [88].

A space X is *strongly Fréchet* at a point $x \in X$ [80] if for each decreasing sequence $\{A_n\}$ of subsets of X with $x \in \bigcap_{n\in\mathbb{N}} \overline{A_n}$, there is an $x_n \in A_n$ for each $n \in \mathbb{N}$ such that the sequence $\{x_n\}$ converges to the point x. A space X is strongly Fréchet [80] if it is strongly Fréchet at each point of X.

Question 3.2. ([43, Question 2.6.19], [55]) Is each almost-open mapping on a strongly Fréchet space a sequence-covering mapping?

Sequence-covering mappings are 1-sequence-covering mappings under certain conditions. For example, boundary-compact sequence-covering mappings on first-countable spaces are 1-sequence-covering mappings [43, Theorem 3.5.3], in particularly, compact sequence-covering mappings on metric spaces are 1-sequence-covering mappings [51].

A regular space X is called *g-metrizable* [81] if X has a σ-locally finite weak-base.

Question 3.3. ([43, Question 3.5.8]) Let X and Y be topological spaces, and $f : X \to Y$ a boundary-compact sequence-covering mapping. Is f a 1-sequence-covering mapping if X satisfies one of the following conditions?

(1) every compact subset of X has a countable sn-network in X.
(2) every compact subset of X has a countable outer sn-network in X.
(3) X has a compact-countable sn-network.
(4) X is a g-metrizable space.

The following relations hold in the conditions above: (4) $\Rightarrow$ (3) $\Rightarrow$ (2) and (1).

Spaces with a point-countable base are preserved by countably bi-quotient s-mappings [18]. What kind of mappings can preserve spaces with a point-countable sn-network (resp. weak-base)? Spaces with a point-countable sn-network are preserved by countable-to-one 1-sequence-covering mappings [34]. Moreover, spaces with a point-countable sn-network can be characterized as 1-sequence-covering, at most boundary-one and s-images of metric spaces [43, Theorem 2.6.12].

Question 3.4. ([43, Question 2.6.21], [34]) Is every space with a point-countable sn-network preserved by 1-sequence-covering, at most boundary-one and s-mappings?

Sequence-covering and closed images of regular spaces with a point-countable weak-base are gf-countable [57].

Question 3.5. ([43, Question 3.5.19], [35]) Let $f : X \to Y$ be a sequence-covering and closed mapping. Is Y an snf-countable space if X has a point-countable sn-network?

Sequence-covering and closed images of regular spaces with a σ-compact-finite weak-base also have a σ-compact-finite weak-base [57].

Question 3.6. ([43, Question 4.1.29], [37]) Does every sequence-covering and closed image of a space with a σ-compact-finite sn-network have a σ-compact-finite sn-network?

Regular spaces with a point-countable base are preserved by sequence-covering and closed mappings [59].

Question 3.7. ([43, Question 3.5.18]) Let $f : X \to Y$ be a sequence-covering and closed mapping. Does Y have a point-countable weak-base if X is a regular space with a point-countable weak-base?

We shall strengthen the forms of compact-covering and sequence-covering (resp. 1-sequence-covering) mappings. Let $f : X \to Y$ be a mapping. f is called a 1-*scc-mapping* [54] if for each compact subset $K \subset Y$, there is a compact subset $L \subset X$ such that $f(L) = K$ and for each $y \in K$, there is a point $x \in L$ such that if whenever $\{y_n\}$ is a sequence converging to y, there exists a sequence $\{x_n\}$ converging to x in X with each $x_n \in f^{-1}(y_n)$. f is called an *scc-mapping* [54] if for each compact subset $K \subset Y$, there is a compact subset $L \subset X$ such that $f(L) = K$ and if whenever $\{y_n\}$ is a sequence converging to some point in K, there exists a sequence $\{x_n\}$ converging to some point in L with each $x_n \in f^{-1}(y_n)$.

The following are known that

(1) Every 1-*scc*-mapping (resp. *scc*-mapping) is a 1-sequence-covering (resp. sequence-covering) and compact-covering mapping.
(2) Every compact-covering open mapping on a first-countable space is a 1-*scc*-mapping [54].
(3) Every *scc*-mapping on a first-countable space is a 1-*scc*-mapping [43, Theorem 2.7.15].
(4) There exists a 1-sequence-covering compact-covering mapping on a metric space which is not an *scc*-mapping [54].

Question 3.8. ([43, Question 2.7.16], [54]) Is every *scc*-mapping on compact spaces a 1-*scc*-mapping?

Sequence-covering mappings are pseudo-sequence-covering mappings and sequentially quotient mappings. For a topological space X every sequence-covering mapping onto the space X is a 1-sequence-covering mapping if and only if for each point $x \in X$, there is a sequence $\{x_n\}$ converging to x in X such that the set $\{x_n : n \in \mathbb{N}\} \cup \{x\}$ is a sequential neighborhood of x in X [46]. On the other hand, every sequentially quotient mapping onto a space X is a 1-sequence-covering mapping if and only if there is a nontrivial convergent sequence in X [46].

Question 3.9. ([43, Question 3.5.27]) Give a characterization of a space X such that every pseudo-sequence-covering mapping onto the space X is a 1-sequence-covering mapping.

4. Images of metric spaces

In the past fifty years, many interesting works are induced by certain mappings on metric spaces. Obviously, every pseudo-sequence-covering mapping on a metric space is a sequentially quotient mapping [43, Lemma 1.3.4]. Every sequentially quotient and boundary-compact mapping on a developable space or a space with a point-countable base is a pseudo-sequence-covering mapping [36].

Question 4.1. ([43, Question 3.5.21]) Is every sequentially quotient s-mapping on a metric space a pseudo-sequence-covering mapping?

Every snf-countable space can be characterized as a sequentially quotient and boundary-compact image of a metric space [34]. The sequentially quotient, at most boundary-one and s-image of a metric space can be characterized as a space with a point-countable sn-network [34].

Question 4.2. ([43, Question 2.6.22], [34]) Let X be a sequentially quotient s-image of a metric space. Is X a sequentially quotient, boundary-compact and s-image of a metric space if X is snf-countable?

A topological space X is called *feebly compact* [82] if every locally finite family of open subsets of X is finite. It is clear that countably compact

spaces are feebly compact, and that feebly compact spaces are pseudo-compact. Arhangel'skiĭ [9] showed that if a regular feebly compact space X is a pseudo-open s-image of a metric space, then X has a point-countable base. If a countably compact space X is a quotient s-image of a metric space, then X is metrizable [23]. Here, a mapping $f : X \to Y$ is called *pseudo-open* [7] if V is an open subset of X and $f^{-1}(y) \subset V$ for some $y \in Y$, then $f(V)$ is a neighborhood of y in Y. Obviously, every pseudo-open mapping is a quotient mapping.

Question 4.3. ([43, Question 2.2.12]) Let X be a feebly compact space. Does X have a point-countable base if X is a quotient s-image of a metric space?

Liu [58] proved that a space is a quotient, at most boundary-one and s-image of a metric space if and only if it is a quotient, at most boundary-one and countable-to-one image of a metric space.

Question 4.4. ([43, Question 3.4.14], [34]) Is every quotient compact image of a metric space a countable-to-one quotient image of a metric space?

While locally separable metric spaces are between separable metric spaces and metric spaces, the images of locally separable metric spaces are very different from the images of separable metric spaces, or the images of metric spaces [43, Example 5.1.23]. Sequentially quotient s-images of metric spaces are equivalent to pseudo-sequence-covering s-images of metric spaces.

Question 4.5. ([43, Question 5.1.8]) Is every sequentially quotient s-image of a locally separable metric space equivalent to a pseudo-sequence-covering s-image of a locally separable metric space?

Although some intrinsic characterizations about quotient s-images of locally separable metric spaces have been obtained, most of these characterizations are more complicated [43, Theorems 5.1.4, 5.1.7 and 5.1.9]. It's still an open question: how to seek a nice intrinsic characterization of a quotient s-image of a locally separable metric space [62, 84]. Some attempts have been made to solve the above question by the following questions.

Question 4.6. ([43, Question 5.1.15], [44]) Is each closed Lindelöf subspace of a space X is separable if the space X is a quotient s-image of a locally separable metric space?

Every first-countable subspace of a quotient s-image of a locally separable metric space is locally separable. However, there exists a space X which is a quotient s-image of a metric space such that every first-countable subspace of the space X is locally separable, but X is not any quotient s-image of a locally separable metric space [48, 75].

Question 4.7. ([43, Question 5.1.16], [44]) Let X be a quotient s-image of a metric space. If every first-countable subspace of X is locally separable and every closed Lindelöf subspace of X is separable, is X a quotient s-image of a locally separable metric space?

A topological space X is called a *cosmic space* (resp. an $\aleph_0$-*space*) [66] if it has a countable network (resp. k-network). Cosmic spaces and $\aleph_0$-spaces are very nice spaces represented as certain images of metric spaces. Michael [66] proved that a space is a cosmic space if and only if it is an image of separable metric space, and that a space is an $\aleph_0$-space if and only if it is a compact-covering image of separable metric space.

It is clear that every quotient s-image of a locally separable metric space is a sequential space with a point-countable cs^*-network consisting of cosmic subspaces.

Question 4.8. ([43, Question 5.1.24], [44]) Is every sequential space with a point-countable cs^*-network consisting of cosmic subspaces a quotient s-image of a locally separable metric space?

Let $\mathscr{P}$ be a cover of a topological space X. $\mathscr{P}$ is called a cs^*-*cover* of the space X [31] if, for each convergent sequence S in X, there exists a $P \in \mathscr{P}$ such that some subsequence of S is frequently in P. $\mathscr{P}$ is called a k-*cover* of the space X [65] if every compact subset of X is covered by some finite subfamily of $\mathscr{P}$. Spaces with a point-countable cs^*-cover (resp. k-cover) consisting of $\aleph_0$-subspaces are pseudo-sequence-covering (resp. compact-covering) s-images of locally separable metric spaces.

Question 4.9. ([43, Question 5.1.25], [32, 33])

(1) Is every pseudo-sequence-covering s-image of locally separable metric spaces a space with a point-countable cs^*-cover consisting of $\aleph_0$-subspaces? It is even unknown whether every quotient s-image of locally separable metric spaces has a point-countable cs^*-network consisting of $\aleph_0$-subspaces [47, 62].
(2) Is every compact-covering s-image of locally separable metric spaces a space with a point-countable k-cover consisting of $\aleph_0$-subspaces?

Sequentially quotient compact images of locally separable metric spaces are equivalent to pseudo-sequence-covering compact images of locally separable metric spaces [20]. A characterization of the images has been obtained, but it is quite complex [64]. Let X be a topological space and $\{\mathscr{P}_n\}$ a sequence of covers of the space X. $\{\mathscr{P}_n\}$ is called a *point-star network* [52] for X if the family $\{\mathrm{st}(x, \mathscr{P}_n) : n \in \mathbb{N}\}$ is a network of x in X for each $x \in X$.

Question 4.10. ([43, Question 5.2.4])

(1) Let X be a space with a point-star network of point-finite cs^*-covers consisting of cosmic subspaces. Is X a pseudo-sequence-covering compact image of a locally separable metric space [44]?
(2) Let X be a sequential space with a point-star network of point-finite cs^*-covers consisting of cosmic subspaces. Is X a quotient compact image of a locally separable metric space [21]?

Let (X, d) be a metric space and Y a topological space. A mapping $f : X \to Y$ is called a π-*mapping* [72] if $d(f^{-1}(y), X \setminus f^{-1}(U)) > 0$ for each

$y \in Y$ and each neighborhood U of y in Y. Each compact mapping is a π-mapping on a metric space.

Question 4.11. ([43, Question 5.2.11], [2]) Is every quotient π-image of a locally separable metric space equivalent to a pseudo-sequence-covering quotient π-image of a locally separable metric space?

Quotient compact images of locally separable metric spaces are not necessarily quotient compact and compact-covering images of metric spaces [15]. Quotient π-images of separable metric spaces are pseudo-sequence-covering quotient and compact images of separable metric spaces [4]. Quotient compact and regular images of separable metric spaces are quotient compact and compact-covering images of separable metric spaces [51].

Question 4.12. ([43, Question 3.3.24]) Is every quotient compact image of a separable metric space a quotient compact and compact-covering image of a separable metric space?

5. Hereditarily closure-preserving families

The research about hereditarily closure-preserving families is mainly originated from closed images of locally finite families in a topological space. Let $\mathscr{P}$ be a family of subsets of a topological space X. $\mathscr{P}$ is called a *hereditarily closure-preserving family* [30] of the space X if the family $\{H(P) : P \in \mathscr{P}\}$ is closure-preserved for each $H(P) \subset P \in \mathscr{P}$, i.e.,

$$\overline{\cup\{H(P) : P \in \mathscr{P}'\}} = \cup\{\overline{H(P)} : P \in \mathscr{P}'\}$$

for each $\mathscr{P}' \subset \mathscr{P}$. $\mathscr{P}$ is called a *point-discrete family* [60] or a *weakly hereditarily closure-preserving family* [14] of the space X if the family $\{\{p(P)\} : P \in \mathscr{P}\}$ is closure-preserving for each $p(P) \in P \in \mathscr{P}$, i.e., the set $\{p(P) : P \in \mathscr{P}\}$ is closed discrete in X. Obviously, every locally finite family is hereditarily closure-preserving, and every hereditarily closure-preserving family is point-discrete.

In regular spaces, the closure $\overline{\mathscr{P}} = \{\overline{P} : P \in \mathscr{P}\}$ of a hereditarily closure-preserving family $\mathscr{P}$ is still hereditarily closure-preserving [38, 68]. Generally, we know the closure of a hereditarily closure-preserving family is point-discrete [77].

Question 5.1. ([43, Question 4.2.7], [41]) Is the closure of every hereditarily closure-preserving family still hereditarily closure-preserving?

About the relations between a σ-point-discrete family and a σ-compact-finite family, it is known that a space with a σ-point-discrete sn-network (resp. k-network, wcs^*-network, network) has a σ-compact-finite sn-network [37](resp. k-network [73], wcs^*-network [78], network [43]). However, there exists a space with a σ-point-discrete base (resp. weak-base) which does not have a σ-compact-finite base [14] (resp. weak-base [50]).

Question 5.2. ([43, Question 4.1.19]) Does every space with a σ-point-discrete cs-network have a σ-compact-finite cs-network [78] or a σ-compact-finite cs^*-network?

Sometimes, an snf-countable space having certain cs-networks may imply that it has certain sn-networks. For example, a space with a point-countable sn-network if and only if it is an snf-countable space with a point-countable cs-network [53].

Question 5.3. ([43, Question 4.1.20]) Does every snf-countable space with a σ-point-discrete cs-network have a σ-point-discrete sn-network?

Because an snf-countable space with a σ-point-discrete cs-network has a σ-compact-finite sn-network [35, 37], a gf-countable space with a σ-point-discrete cs-network has a σ-compact-finite weak-base. However, there exists a gf-countable space with a σ-point-discrete wcs^*-network which is not a space having a σ-point-discrete cs^*-network [43, Example 4.1.16(7)].

Question 5.4. ([43, Question 4.1.22], [78]) Does every gf-countable space with a σ-point-discrete cs^*-network have a σ-compact-finite weak-base?

Liu and Ludwig [61] proved each closed mapping on a regular space with a σ-point-discrete base is a compact-covering mapping. Liu, Lin and Ludwig [60] proved that every closed mapping on a regular space with a σ-point-discrete weak-base is also a compact-covering mapping under (CH).

Question 5.5. ([43, Question 4.1.27]) Is every closed mapping on a regular space with a σ-point-discrete sn-network a compact-covering mapping?

Question 5.6. ([43, Question 4.1.28], [37]) Is every space with a σ-point-discrete sn-network preserved by a sequence-covering closed mapping?

Lin, Liu and Dai [47] proved that a regular space X with a σ-hereditarily closure-preserving k-network consisting of $\aleph_0$-subspaces if and only if X has a σ-hereditarily closure-preserving k-network and each first-countable closed subspace of X is locally separable.

Question 5.7. ([43, Question 4.3.11]) Is each first-countable subset of a regular space with a σ-hereditarily closure-preserving k-network consisting of separable subspaces locally separable?

References

[1] P. S. Alexandroff, *On the metrization of topological spaces* (in Russian), Bull. Acad. Pol. Sci., Sér. Sci. Math. Astron. Phys. **8**(1960), 135–140.

[2] T. V. An, N. V. Dung, *On π-images of locally separable metric spaces*, Inter. J. Math. Math. Sci. 2008, Article ID 145683, 8 pages, doi: 10.1155/2008/145683.

[3] T. V. An, L. Q. Tuyen, *On an affirmative answer to S. Lin's problem*, Topology Appl., **158**(2011), 1567–1570.

[4] T. V. An, L. Q. Tuyen, *On π-images of separable metric spaces and a problem of Shou Lin*, Mat. Vesnik. **64**(2012), no. 4, 297–302.

[5] A. V. Arhangel'skiĭ, *On mappings of metric spaces* (in Russian), Dokl. Akad. Nauk SSSR **145**(1962), no. 2, 245–247.

[6] A. V. Arhangel'skiĭ, *On open and almost open mappings of topological spaces* (in Russian), Dokl. Akad. Nauk SSSR **147**(1962), 999–1002.

[7] A. V. Arhangel'skiĭ, *Some types of factor mappings and the relations between classes of topological spaces* (in Russian), Dokl. Akad. Nauk SSSR **153**(1963), 743–746.

[8] A. V. Arhangel'skiĭ, *Mappings and spaces* (in Russian), Uspechi Mat. Nauk **21**(1966), no. 4, 133–184.

[9] A. V. Arhangel'skiĭ, *Components of first-countability and various kinds of pseudoopen mappings*, Topology Appl. **158**(2011), 215–222.

[10] T. Banakh, V. Bogachev and A. Kolesnikov, *k^*-metrizable spaces and their applications*, J. Math. Sci. **155**(2008), no. 4, 475–522.

[11] S. Barov, *A note on the spaces which are quotient compact-covering s-images of metric spaces*, C. R. Acad. Bulgare. Sci. **52**(1999), no. 5-6, 11–14.

[12] J. R. Boone, *Some characterizations of paracompactness in k-spaces*, Fund. Math. **72**(1971), 145–153.

[13] J. R. Boone, F. Siwiec, *Sequentially quotient mappings*, Czech. Math. J. **26**(1976), 174–182.

[14] D. K. Burke, R. Engelking and D. J. Lutzer, *Hereditarily closure-preserving collections and metrization*, Proc. Amer. Math. Soc. **51**(1975), no. 2, 483–488.

[15] Huaipeng Chen, *Weak neighborhoods and Michael-Nagami's question*, Houston J. Math. **25**(1999), no. 2, 297–309.

[16] Huaipeng Chen, *Compact-covering maps and k-networks*, Proc. Amer. Math. Soc. **131**(2003), no. 8, 2623–2632.

[17] R. Engelking, *General Topology* (revised and completed edition), Berlin: Heldermann Verlag, 1989.

[18] V. V. Filippov, *Quotient spaces and multiplicity of a base* (in Russian), Mat. Sb. **80**(1969), no. 4, 521–532.

[19] Zhimin Gao, *$\aleph$-space is invariant under perfect mappings*, Questions Answers in General Topology **5**(1987), no. 2, 271–279.

[20] Ying Ge, *On compact images of locally separable metric spaces*, Topology Proc. **27**(2003), no. 1, 351–360.

[21] Ying Ge, Shou Lin, *Uniform covers and compact images of metric spaces* (in Chinese), Acta Math. Sinica **47**(2004), no. 6, 1149–1154.

[22] G. Gruenhage, *Generalized metric spaces*, In: K. Kunen, J. E. Vaughan eds., Handbook of Set-theoretic Topology, Amsterdam: North-Holland, 1984, 423–501.

[23] G. Gruenhage, E. A. Michael and Y. Tanaka, *Spaces determined by point-countable covers*, Pacific J. Math. **113**(1984), no. 2, 303–332.

[24] J. A. Guthrie, *A characterization of $\aleph_0$-spaces*, General Topology Appl. **1**(1971), 105–110.

[25] K. P. Hart, J. van Mill and P. Simon, *Recent Progress in General Topology III*, Papers from the Prague Topological Symp., Prague, 2011. Amsterdam: Elsevier Science Publishers B. V., 2014.

[26] K. P. Hart, J. Nagata and J. E. Vaughan, *Encyclopedia of General Topology*, Amsterdam: Elsevier Science Publishers B. V., 2004.

[27] M. Hušek, J. van Mill, *Recent Progress in General Topology*, Papers from the Prague Topological Symp., Prague, 1991. Amsterdam: Elsevier Science Publishers B. V., 1992.

[28] M. Hušek, J. van Mill. *Recent Progress in General Topology II*, Papers from the Prague Topological Symp. Prague, 2001. Amsterdam: Elsevier Science Publishers B. V., 2002.

[29] Y. Ikeda, Chuan Liu and Y. Tanaka, *Quotient compact images of metric spaces, and related matters*, Topology Appl. **122**(2002), no. 1-2, 237–252.

[30] N. Lašnev, *Closed images of metric spaces* (in Russian), Dokl. Akad. Nauk SSSR **170**(1966), no. 3, 505–507.

[31] Jinjin Li, *Images on locally separable metric spaces and related results*, Doctoral Dissertation, Shandong University, Jinan, 2000.

[32] Jinjin Li, Weiyuan Cai, *A note on sequence-covering s-mappings* (in Chinese), Acta Math. Sinica (Chin. Ser.) **43**(2000), no. 4, 757–762.

[33] Jinjin Li, Shouli Jiang, *Properties of some special covers*, Indian J. Math. **44**(2002), no. 3, 345–351.

[34] Fucai Lin, Shou Lin, *On sequence-covering boundary-compact maps of metric spaces*, Adv. Math. (China) **39**(2010), no. 1, 71–78.

[35] Fucai Lin, Shou Lin, *Some notes on closed sequence-covering maps*, J. Math. Resear. Appl. **34**(2014), no. 1, 97–104.

[36] Fucai Lin, Shou Lin, *Sequence-covering maps on generalized metric spaces*, Houston J. Math. **40**(2014), no. 3, 927–943.

[37] Fucai Lin, Rongxin Shen, *Some notes on σ-point-discrete sn-networks*, Adv. Math. (China) **39**(2010), no. 2, 212–216.

[38] Shou Lin, *On a problem of K. Tamano*, Questions Answers in General Topology **6**(1988), 99–102.

[39] Shou Lin, *On sequence-covering s-mappings* (in Chinese), Adv. Math. (China) **25**(1996), no. 6, 548–551.

[40] Shou Lin, *A note on the Arens' space and sequential fan*, Topology Appl. **81**(1997), no. 3, 185–196.

[41] Shou Lin, *Regularity in "Generalized Metric Spaces and Mappings"*, J. Ningde Teachers College (Natural Sci.) **11**(1999), no. 4, 241–247.

[42] Shou Lin, *Point-countable covers and sequence-covering mappings*, Doctoral Dissertation, Zhejang University, Hangzhou, 2000.

[43] Shou Lin, *Point-countable Covers and Sequence-covering Mappings* (second edition), Beijing: China Science Press, 2015.

[44] Shou Lin, Zhangyong Cai and Jinjin Li, *Recent advances on the images of locally separable metric spaces* (in Chinese), Adv. Math. (China) **38**(2009), no. 6, 657–663.

[45] Shou Lin, Jinjin Li, *Semi-quotient mappings and spaces with compact-countable k-networks*, Adv. Math. (China) **38**(2009), no, 417–426.

[46] Shou Lin, Kedian Li and Ying Ge, *Convergent-sequence spaces and sequence-covering mappings*, Houston J. Math. **39**(2013), no. 4, 1367–1384.

[47] Shou Lin, Chuan Liu and Mumin Dai, *Images on locally separable metric spaces*, Acta Math. Sinica (New Ser.) **13**(1997), no. 1, 1–8.

[48] Shou Lin, M. Sakai, *Counterexamples on the images of locally separable metric spaces*, Topology Proc. **31**(2007), no. 1, 181–187.

[49] Shou Lin, Y Tanaka, *Point-countable k-networks, closed maps, and related results*, Topology Appl. **59**(1994), no. 1, 79–86.

[50] Shou Lin, Li Yan, *A note on spaces with a σ-compact-finite weak base*, Tsukuba J. Math. **28**(2004), 85–91.

[51] Shou Lin, Pengfei Yan, *Sequence-covering maps of metric spaces*, Topology Appl. **109**(2001), no. 3, 301–314.

[52] Shou Lin, Pengfei Yan, *On sequence-covering compact mappings* (in Chinese), Acta Math. Sinica (Chin. Ser.) **44**(2001), no. 1, 175–182.

[53] Shou Lin, Pengfei Yan, *On a theorem of D. Burke and E. Michael* (in Chinese), Chinese Ann. Math. **22A**(2001), no. 6, 707–714.

[54] Shou Lin, Jing Zhang, *Weak bases and the compact-covering images of metric spaces* (in Chinese), Acta Math. Sci. **33A**(2013), no. 3, 483–493.

[55] Shou Lin, Zhongjing Zhu, *Workshop lecture on the images of some metric spaces* (in Chinese), Adv. Math. (China) **42**(2013), no. 2, 129–137.

[56] Chuan Liu, *Spaces with a σ-compact finite k-network*, Questions Answers in General Topology **l0**(1992), 81–87.

[57] Chuan Liu, *On weak bases*, Topology Appl. **150**(2005), 91–99.

[58] Chuan Liu, *A note on point-countable weak bases*, Questions Answers in General Topology **25**(2007), 57–61.

[59] Chuan Liu, *Notes on closed mappings*, Houston J. Math. **33**(2007), no. 1, 249–259.

[60] Chuan Liu, Shou Lin and L. D. Ludwig, *Spaces with a σ-point-discrete weak base*, Tsukuba J. Math. **32**(2008), no. 1, 165–177.

[61] Chuan Liu, L. D. Ludwig, *Nagata-Smirnov revisited: spaces with σ-WHCP bases*, Topology Proc. **29**(2005), no. 2, 559–565.

[62] Chuan Liu, Y. Tanaka, *Spaces with certain compact-countable k-networks, and questions*, Questions Answers in General Topology **14**(1996), no. 1, 15–37.

[63] Chuan Liu, Y. Tanaka, *Spaces and mappings: special networks*, In: E. Pearl ed., Open Problems in Topology II, Amsterdam: Elsevier Science Publishers B. V., 2007, 23–34.

[64] Cheng Lv, Hongyan Li, *Pseudo-sequence-covering images of locally separable metric spaces* (in Chinese), College Math. (China) **21**(2005), no. 3, 52–56.

[65] R. A. McCoy, I. Ntantu, *Topological properties of spaces of continuous functions*, In: Lecture Notes in Math., no. 1315, Berlin: Springer-Verlag, 1988.

[66] E. A. Michael, *$\aleph_0$-spaces*, J. Math. Mech. **15**(1966), no. 6, 983–1002.

[67] E. A. Michael, K. Nagami, *Compact-covering images of metric spaces*, Proc. Amer. Math. Soc. **37**(1973), no. 1, 260–266.

[68] E. A. Micheal, R. C. Olson and F. Siwiec, *A-spaces and countably bi-quotient maps*, Dissertationes Math. **33**(1976), 1–43.

[69] J. van Mill, G. M. Reed. *Open Problems in Topology*, Amsterdam: Elsevier Science Publishers B. V., 1990.

[70] P. O'Meara, *On paracompactness in function spaces with the compact-open topology*, Proc. Amer. Math. Soc. **29**(1971), no. 1, 183–189.

[71] E. Pearl, *Open Problems in Topology II*, Amsterdam: Elsevier Science Publishers B. V., 2007.

[72] V. I. Ponomarev, *Axioms of countability and continuous mappings* (in Russian), Bull. Acad. Pol. Sci., Sér. Sci. Math. Astron. Phys. **8**(1960), 127–134.

[73] Zhibin Qu, Zhimin Gao, *Spaces with compact-countable k-networks*, Math. Japonica. **49**(1999), no. 2, 199–205.

[74] M. Sakai, *Remarks on spaces with special type of k-networks*, Tsukuba J. Math. **21**(1997), no. 2, 443–448.

[75] M. Sakai, *Counterexamples on generalized metric spaces*, Sci. Math. Japonica. **64**(2006), no. 1, 73–76.

[76] D. B. Shakhmatov, *On pseudocompact spaces with point-countable bases*, Soviet Math. Dokl. **30**(1984), 747–751.

[77] Jianhua Shen, Ying Ge and Zhihong Ge, *A note on weakly hereditarily closure-preserving families*, Siberian Elec. Math. Reports **3**(2006), 253–256.

[78] Rongxin Shen, Shou Lin, *Spaces with σ-point-discrete $\aleph_0$-weak bases*, Appl. Math. J. Chinese. Univ. **28B**(2013), no. 1, 116–126.

[79] A. Shibakov, *On spaces with point-countable k-networks and their mappings*, Serdica-Bulgaricae Math. Pub. **20**(1994), no. 1, 48–55.

[80] F. Siwiec, *Sequence-covering and countably bi-quotient mappings*, General Topology Appl. **1**(1971), 143–154.

[81] F. Siwiec, *On defining a space by a weak base*, Pacific J. Math. **52**(1974), no. 1, 233–245.

[82] R. M. Stephenson Jr., *Symmetrizable, $\mathscr{F}$-, and weakly first countable spaces*, Can. J. Math. **29**(1977), 480–488.

[83] Y. Tanaka, *Point-countable covers and k-networks,* Topology Proc. **12**(1987), 327–349.

[84] Y. Tanaka, Shengxiang Xia, *Certain s-images of locally separable metric spaces*, Questions Answers in General Topology **14**(1996), no. 2, 217–231.

[85] V. V. Uspenskii, *Pseudocompact spaces with a σ-point-finite base are metrizable*, Comment. Math. Univ. Carolina. **25**(1984), 261–264.

[86] W. S. Watson, *A pseudocompact meta-Lindelöf space which is not compact*, Topology Appl. **20**(1985), 237–243.

[87] Pengfei Yan, Shou Lin, *The compact-covering s-mappings on metric spaces* (in Chinese), Acta Math. Sinica (Chin. Ser.) **42**(1999), no. 2, 241–244.

[88] Pengfei Yan, Shou Lin and Hairong Yang, *On finite subsequence-covering maps*, Adv. Math. (China) **39**(2010), no. 6, 651–656.

Shou Lin(the corresponding author): Institute of Mathematics, Ningde Normal University, Ningde 352100, PR China;
E-mail address: shoulin60@163.com

Jing Zhang: School of Mathematics and Statistics, Minnan Normal University, Zhangzhou 363000, PR China
E-mail address: zhangjing86@126.com